

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07C 217/48, 213/06, C07B 57/00	A1	(11) International Publication Number: WO 00/61540 (43) International Publication Date: 19 October 2000 (19.10.00)
(21) International Application Number: PCT/US00/06423 (22) International Filing Date: 22 March 2000 (22.03.00) (30) Priority Data: 60/128,480 9 April 1999 (09.04.99) US (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): KJELL, Douglas, Patton [US/US]; 3075 Hamilton Street, West Lafayette, IN 47906 (US). LORENZ, Kurt, Thomas [US/US]; 1405 Castle Drive, Lafayette, IN 47905 (US). (74) Agent: STEMERICK, David, M.; Lilly Corporate Center, Indianapolis, IN 46285 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: METHODS FOR PREPARING 3-ARYLOXY-3-ARYLPROPYLAMINES AND INTERMEDIATES THEREOF (57) Abstract The present invention provides processes for the preparation of 3-aryloxy-3-arylpropylamines and intermediates thereof using an nucleophilic aromatic displacement in 1,3-dimethyl-2-imidazolidinone or N-methylpyrrolidinone.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-

5

10 METHODS FOR PREPARING 3-ARYLOXY-3-ARYLPROPYLAMINES
 AND INTERMEDIATES THEREOF

 This application claims the benefit of US Serial Number
60/128,480, filed April 9, 1999.

15

 Certain 3-aryloxy-3-arylpropylamines are known to
possess central nervous system activity. See U.S. Patent
Nos. 5,744,474; 5,023,269; 4,956,388; 4,194,009; 4,314,081
and 5,658,590. The present application relates to processes
20 for preparing 3-aryloxy-3-arylpropylamines, including
duloxetine and tomoxetine.

 Syntheses of 3-aryloxy-3-arylpropylamines utilizing
nucleophilic aromatic displacement are known in the art.
For example U.S. Patent Nos. 5,225,585; 5,166,437;
25 5,023,269; 4,956,388; 5,362,886; and 5,023,269; Tetrahedron
Letters, 31(49), 7101-7104 (1990); and PCT Publication No.
WO 94/00416.

 The nucleophilic aromatic displacement reaction with 3-
hydroxy-3-arylpropylamines is facile for activated aryl
30 halides. A variety of dipolar solvents, for example,
dimethylsulfoxide (WO 94/00416) and 1,3-dimethyl-2-
imidazolidinone and N-methylpyrrolidinone (US Patent No.
5,847,214) have been reported for the reaction of N-methyl-
3-phenyl-3-hydroxypropylamine with 4-trifluoromethyl-1-
35 chlorobenzene to give N-methyl-(4-trifluoromethylphenoxy)-3-
phenylpropylamine (fluoxetine).

 Nucleophilic aromatic displacement of alkoxides in 1,3-
dimethyl-2-imidazolidinone are described in Japanese Kokoku
Patent Publication Sho 60-23656, published June 8, 1985.
40 However, the description is limited to lower order alcohols

-2-

and the exemplified preparations using unactivated aromatics are carried out at temperatures of from 160° to 190°C in a pressure vessel reactor. See Japanese Kokoku Patent Publication Sho 60-23656, published June 8, 1985, examples 5 7-9, 11, and 14. Thus, it does not appear that 1,3-dimethyl-2-imidazolidinone would be a useful solvent in a safe and convenient nucleophilic aromatic displacement with complex alcohols using unactivated aromatics. This is especially so for aromatics such as 2-fluorotoluene which 10 has a boiling point of 113-114°C.

In addition, the reported reaction of unactivated substrate, 2-fluorotoluene, with the alkoxide of (S)-N-methyl-3-phenyl-3-hydroxypropylamine in dimethylsulfoxide gave a modest yield. Tet. Let., 35, 1339-1342 (1994).


15 In spite of the difficulties with using unactivated aryl halides, a method for preparing 3-aryloxy-3-arylpropylamines using nucleophilic aromatic displacement is desirable. In contrast to methods utilizing other displacements, such as the Mitsunobu reaction or 20 displacement of a halide by a phenol, the nucleophilic aromatic displacement method allows for cost efficient assembly of the required substituents directly from a 3-hydroxy-3-arylpropylamine.

Surprisingly, we have discovered that nucleophilic 25 aromatic displacement using complex benzylic alcohols, such as N-methyl-3-phenyl-3-hydroxypropylamine and N,N-dimethyl-3-(2-thienyl)-3-hydroxypropylamine can be carried out with unactivated aromatics, such as 1-fluoronaphthylene and 2-fluorotoluene in 1,3-dimethyl-2-imidazolidinone or N-methylpyrrolidinone at temperatures of less than about 30 140°C.

The present processes provide safe and convenient methods for a high yield preparation of 3-aryloxy-3-arylpropylamines utilizing nucleophilic aromatic 35 displacement on unactivated aromatics in 1,3-dimethyl-2-imidazolidinone or N-methylpyrrolidinone.

-3-

The present invention relates to a process for preparing a 3-aryloxy-3-arylpropylamine of the formula



5

wherein

Ar is phenyl or 2-thienyl,

Ar₁ is 1-naphthyl, 2-methoxyphenyl, 2-thiomethoxyphenyl, or 2-methylphenyl;


10

G is hydrogen or methyl,

and the pharmaceutically-acceptable addition salts thereof comprising the steps of:

(a) reacting an alkoxide of a 3-hydroxy-3-arylpropylamine of the formula

15



wherein

Ar and G are as defined above

20

with a haloaromatic of the formula



wherein

Ar₁-X is 2-fluorotoluene, 2-chlorotoluene, 1-

fluoronaphthalene, 1-chloronaphthalene, 2-fluoroanisole, 2-

25

chloroanisole, 2-fluorothioanisole, or 2-chlorothioanisole,

in 1,3-dimethyl-2-imidazolidinone or N-methylpyrrolidinone

to give the 3-aryloxy-3-arylpropylamine;

(b) optional N-demethylated of the 3-aryloxy-3-

arylpropylamine wherein G is methyl to give the 3-aryloxy-3-

30


arylpropylamine wherein G is hydrogen;

-4-

(c) optional resolution of the 3-aryloxy-3-arylpropylamine to give a specific isomer of the 3-aryloxy-3-arylpropylamine; and

(d) optional formation of an acid addition salt using a pharmaceutically-acceptable acid.

That is, the present invention provides a process for preparing a 3-aryloxy-3-arylpropylamine, as defined above and the pharmaceutically acceptable salts thereof, which comprises reacting an alkoxide of 3-hydroxy-3-arylpropylamine of the formula



wherein

Ar and G are as defined above with a haloaromatic as defined above, characterized in that, 1,3-dimethyl-2-imidazolidinone or N-methylpyrrolidinone is used as solvent.

Particularly, the present invention relates to a process for preparing tomoxetine and the pharmaceutically-acceptable addition salts thereof comprising the steps of:

- (a) reacting an alkoxide of N-methyl-3-phenyl-3-hydroxypropylamine with 2-fluorotoluene in 1,3-dimethyl-2-imidazolidinone to give N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine;
- (b) resolution of N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine to give (R)-N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine (tomoxetine); and
- (c) optional formation of an acid addition salt using a pharmaceutically-acceptable acid.

The present invention also relates to a particular process for preparing tomoxetine and the pharmaceutically-acceptable addition salts thereof comprising the steps of:

- (a) reacting an alkoxide of N,N-dimethyl-3-phenyl-3-hydroxypropylamine) with 2-fluorotoluene in 1,3-dimethyl-2-

-5-

imidazolidinone to give N,N-dimethyl-3-(2-methylphenoxy)-3-phenylpropylamine;

(b) N-demethylated N,N-dimethyl-3-(2-methylphenoxy)-3-phenylpropylamine to give N-methyl-3-(2-methylphenoxy)-3-

5 phenylpropylamine;

(c) resolution of N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine to give (R)-N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine (tomoxetine); and

(d) optional formation of an acid addition salt using a
10 pharmaceutically-acceptable acid.

That is, the present invention relates to processes for preparing tomoxetine and the pharmaceutically-acceptable addition salts thereof, which comprises, reacting an alkoxide of N-methyl-3-phenyl-3-hydroxypropylamine, or an N-
15 protected derivative thereof, with 2-fluorotoluene, characterized in that, 1,3-dimethyl-2-imidazolidinone is used as solvent.

As used herein, the following terms have the meanings indicated:

20 (a) the term "DMI" refers to 1,3-dimethyl-2-imidazolidinone;

(b) the term "NMP" refers to N-methylpyrrolidinone;

(c) the term "ee" or "enantiomeric excess" refers to the percent by which one enantiomer, E_1 , is in excess in a mixture of both enantiomers ($E_1 + E_2$), as calculated by the
25 equation $\{(E_1 - E_2) / (E_1 + E_2)\} \times 100\% = ee$;

(d) the term "pharmaceutically-acceptable addition salt" refers to an acid addition salt using a pharmaceutically-acceptable acid.

The 3-aryloxy-3-arylpropylamines and the intermediates
30 described herein form pharmaceutically acceptable acid addition salts with a wide variety of organic and inorganic acids and include the physiologically acceptable salts which are often used in pharmaceutical chemistry.

A pharmaceutically-acceptable addition salt is formed
35 from a pharmaceutically-acceptable acid as is well known in the art. Such salts are also part of this invention. Typical inorganic acids used to form such salts include

-6-

hydrochloric, hydrobromic, hydriodic, nitric, sulfuric, phosphoric, hypophosphoric, metaphosphoric, pyrophosphoric, and the like. Salts derived from organic acids, such as aliphatic mono and dicarboxylic acids, phenyl substituted alkanolic acids, hydroxyalkanoic and hydroxyalkandioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, may also be used. Such pharmaceutically acceptable salts thus include acetate, phenylacetate, trifluoroacetate, acrylate, ascorbate, benzoate, chlorobenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, methylbenzoate, o-acetoxybenzoate, naphthalene-2-benzoate, bromide, isobutyrate, phenylbutyrate, α -hydroxybutyrate, butyne-1,4-dicarboxylate, hexyne-1,4-dicarboxylate, caprate, caprylate, cinnamate, citrate, formate, fumarate, glycollate, heptanoate, hippurate, lactate, malate, maleate, hydroxymaleate, malonate, mandelate, mesylate, nicotinate, isonicotinate, nitrate, oxalate, phthalate, teraphthalate, propiolate, propionate, phenylpropionate, salicylate, sebacate, succinate, suberate, benzene-sulfonate, p-bromobenzenesulfonate, chlorobenzenesulfonate, ethanesulfonate, 2-hydroxyethanesulfonate, methanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, p-toluenesulfonate, xylenesulfonate, tartarate, and the like.

The present invention relates to processes for the preparation of 3-aryloxy-3-arylpropylamines. It is understood by the skilled person that these compounds exist as stereoisomers. Herein, the Cahn-Prelog-Ingold designations of (R)- and (S)- are used to refer to specific isomers where designated. Specifically, present invention relates to processes for the preparation of duloxetine, (S)-N-methyl-3-(1-naphthyloxy)-3-(2-thienyl)propylamine; and tomoxetine, (R)-N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine. As will be appreciated by the skilled artisan, the present processes are not necessarily limited to the preparation of racemic mixtures or specific isomers. Rather the present processes are capable of preparing both specific enantiomers and racemic mixtures.


-7-

The specific isomers can be obtained by resolution of the product, intermediates, or in some cases the starting materials. For example, duloxetine specific isomers can be most conveniently obtained by utilizing enantiomerically pure starting materials, specifically, (S)-N,N-dimethyl-3-(2-thienyl)-3-hydroxypropylamine or (R)-N-methyl-3-phenyl-3-hydroxypropylamine. As used herein the term "enantiomerically pure" refers to an enatiomeric excess which is greater than 90%, preferably greater than 93%, more preferably greater than 95%.

The present preparation of 3-aryloxy-3-arylpropylamines are carried out according to Reaction Scheme A below. In Reaction Scheme A, all substituents, unless otherwise indicated, are as previously defined. In Reaction Scheme A all reagents are well known and appreciated in the art.

- 8 -

Reaction Scheme A



In Reaction Scheme A, step a, a nucleophilic aromatic displacement, the alkoxide of a suitable 3-hydroxy-3-arylpropylamine of formula (a) is contacted with a suitable haloaromatic, $\text{Ar}_1\text{-X}$, in 1,3-dimethyl-2-imidazolidinone or N-methylpyrrolidinone to give a 3-aryloxy-3-arylpropylamine. As will be appreciated by those skilled in the art, a 3-hydroxy-3-arylpropylamine of formula (a) in which G is methyl gives a 3-aryloxy-3-arylpropylamine of formula (b); a 3-hydroxy-3-arylpropylamine of formula (a) in which G is hydrogen gives a 3-aryloxy-3-arylpropylamine of formula (c); and a specific isomer of a 3-hydroxy-3-arylpropylamine of formula (a) in which G is hydrogen gives directly a 3-aryloxy-3-arylpropylamine of formula (d). Also, it will be appreciated by those skilled in the art that the specific isomers of formula (d) can be obtained by demethylation of a

-9-

3-aryloxy-3-arylpropylamine of formula (b) prepared from a single isomer of a compound of formula (a) in which G is methyl.

Suitable 3-hydroxy-3-arylpropylamines of formula (a) include N,N-dimethyl-3-phenyl-3-hydroxypropylamine, N-methyl-3-phenyl-3-hydroxypropylamine, N,N-dimethyl-3-(2-thienyl)-3-hydroxypropylamine, (R)-N,N-dimethyl-3-phenyl-3-hydroxypropylamine, (R)-N-methyl-3-phenyl-3-hydroxypropylamine, (R)-N,N-dimethyl-3-(2-thienyl)-3-hydroxypropylamine, (S)-N,N-dimethyl-3-phenyl-3-hydroxypropylamine, (S)-N-methyl-3-phenyl-3-hydroxypropylamine, and (S)-N,N-dimethyl-3-(2-thienyl)-3-hydroxypropylamine. Suitable haloaromatics, include 2-fluorotoluene, 2-chlorotoluene, 1-fluoronaphthalene, 1-chloronaphthalene, 2-fluoroanisole, 2-chloroanisole, 2-fluorothioanisole, and 2-chlorothioanisole.

For example, the reaction is carried out using an alkoxide of an alcohol of formula (a). While many metals are suitable for this reaction, generally, an alkali metal alkoxide is used, with the lithium, sodium, and potassium alkoxide being preferred. Sodium and potassium alkoxide are more preferred. The alkoxide is formed by contacting of an alcohol of formula (a) with a suitable base, such as lithium hydride, lithium N,N-diisopropylamide, sodium hydride, potassium hydride, sodium hydroxide, potassium hydroxide, sodium amide, potassium amide, sodium alkoxides, such as sodium t-butoxide, sodium methoxide, sodium ethoxide, potassium alkoxides, such as potassium t-butoxide, potassium methoxide, potassium ethoxide, and the like. From 1 molar equivalent to a large molar excess of base is used. In some cases, when sodium bases are used, the addition of a catalytic amount of potassium ion can advantageously be added, for example, in the form of potassium benzoate.

The reaction is carried out in 1,3-dimethyl-2-imidazolidinone or N-methylpyrrolidinone. Neither high temperatures nor pressure vessels are required. The reaction is carried out at temperatures of from about 0°C to

-10-

about 140°C with temperatures of from about 20°C to about 130°C being preferred, and temperatures of from about 80°C to about 120°C being more preferred. The reaction typically requires from about 1 hour to about 48 hours. The product
5 can be isolated and purified by techniques well known in the art, such as filtration, evaporation, extraction, trituration, chromatography, and crystallization. Alternately, the compound of formula (b) can be taken directly to step c, if desired, without isolation or without
10 purification.

In Reaction Scheme A, optional step b, the compound of the formula (b) is N-demethylated to give the compound of formula (c). As is understood from Reaction Scheme A, where a specific isomer of compound (a) is used to provide a
15 specific isomer of compound (b) the isomer can be N-demethylated to give compound (d) directly. Such N-demethylation reactions are well known and appreciated in the art and include demethylations which proceed through a N-cyano and carbamate intermediates followed by hydrolysis.
20 See for example, U.S. Patent Nos. 4,956,388; 4,314,081; and 5,362,886.

For example, a compound of formula (b) is contacted with a slight molar excess of chloro formate, such as phenyl chloroformate, ethyl chloroformate, trichloroethyl
25 chloroformate, and the like. The reaction is carried out in the presence of a suitable base, such as triethylamine, pyridine, N,N-diisopropylethylamine, and the like. The reaction is carried out in a suitable solvent, such as toluene, dichloromethane, tetrahydrofuran, and the like.
30 Typically the reaction is carried out at temperatures of from about 0°C to the refluxing temperature of the solvent and require about 1 hour to 48 hours. The carbamate intermediate can be isolated and purified by techniques well known in the art, such as filtration, evaporation,
35 extraction, trituration, chromatography, and crystallization. The carbamate intermediate is then

-11-

hydrolyzed to give product. For example, the carbamate intermediate is contacted with an excess of sodium hydroxide or potassium hydroxide. The reaction is carried out in a suitable solvent, such as water, dimethyl sulfoxide, ethanol, dimethyl sulfoxide/water mixtures, and the like. Typically the reaction is carried out at temperatures of from about 20°C to about 100°C and require about 1 hour to 48 hours. The product can be isolated and purified by techniques well known in the art, such as filtration, evaporation, extraction, trituration, chromatography, and crystallization.


In Reaction Scheme A, optional step c, a compound of formula (c) is resolved to give a 3-aryloxy-3-arylpropylamine having a specific stereochemistry. Such resolutions are well known and appreciated in the art, such as the use of mandelic acid as described in European Patent Application No. 0 052 492, published May 26, 1982, the disclosure of which is incorporated by reference. As is understood from Reaction Scheme A, a compound (b) can be resolved by the same techniques to give a specific isomer of compound (b) the isomer can be N-demethylated, if desired to give compound (d) directly.

In Reaction Scheme A, optional step d, not shown, an acid addition salt is formed using a pharmaceutically-acceptable acid. As is appreciated by the person skilled in the art, an acid addition salt can be formed for the end products of formula (b), (c), and (d). The formation of acid addition salts is well known and appreciated in the art.

In particular, the present preparations of tomoxetine are carried out according to Reaction Scheme B below. In Reaction Scheme B, all substituents, unless otherwise indicated, are as previously defined. In Reaction Scheme B all reagents are well known and appreciated in the art.

-12-

Reaction Scheme B



In Reaction Scheme B, step a, a nucleophilic aromatic displacement, the alkoxide of an alcohol of formula (1) in which G is methyl or hydrogen, N,N-dimethyl-3-phenyl-3-hydroxypropylamine where G is methyl or N-methyl-3-phenyl-3-hydroxypropylamine where G is hydrogen, is contacted with a 2-fluorotoluene (the compound of formula (2)) in 1,3-dimethyl-2-imidazolidinone or N-methylpyrrolidinone to give the compound of formula (3), N,N-dimethyl-3-(2-methylphenoxy)-3-phenylpropylamine, or the compound of formula (4), N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine; respectively.

For example, the reaction is carried out using an alkoxide of an alcohol of formula (1). While many metals

-13-

are suitable for this reaction, generally, an alkali metal alkoxide is used, with the lithium, sodium, and potassium alkoxide being preferred. Sodium and potassium alkoxide are more preferred. The alkoxide is formed by contacting of an alcohol of formula (1) with a suitable base, such as lithium hydride, lithium N,N-diisopropylamide, sodium hydride, potassium hydride, sodium amide, potassium amide, sodium alkoxides, such as sodium t-butoxide, potassium alkoxides, such as potassium t-butoxide, and the like. From 1 molar equivalent to a large molar excess of base is used, with about 1.05 to about 1.5 molar equivalents being preferred. The reaction is carried out in 1,3-dimethyl-2-imidazolidinone or N-methylpyrrolidinone with 1,3-dimethyl-2-imidazolidinone being preferred. Neither high temperatures nor pressure apparatus are required. The reaction is carried out at temperatures of from about 75°C to about 140°C with temperatures of from about 90°C to about 130°C being preferred and about 95°C to about 115°C being more preferred. The reaction typically requires from about 12 hours to about 48 hours. The product can be isolated and purified by techniques well known in the art, such as filtration, evaporation, extraction, trituration, chromatography, and crystallization. Alternately, the compound of formula (4) can be taken directly to step c without isolation or without purification.

In Reaction Scheme B, step b, the compound of the formula (3), N,N-dimethyl-3-(2-methylphenoxy)-3-phenylpropylamine, is N-demethylated to give the compound of formula (4), N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine. Such N-demethylation reactions are well known and appreciated in the art and include demethylations which proceed through a N-cyano and carbamate intermediates followed by hydrolysis as described above in Reaction Scheme A, step c, above.

In Reaction Scheme B, step c, N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine is resolved to give(R)-N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine (tomoxetine).

-14-

Such resolutions are well known and appreciated in the art and is described in European Patent Application No. 0 052 492, published May 26, 1982, the disclosure of which is incorporated by reference.

5 In Reaction Scheme B, optional step d, not shown, an acid addition salt is formed using a pharmaceutically-acceptable acid. The formation of acid addition salts is well known and appreciated in the art.

10 The present invention is further illustrated by the following examples and preparations. These examples and preparations are illustrative only and are not intended to limit the invention in any way.

The terms used in the examples and preparations have their normal meanings unless otherwise designated. For
15 example "°C" refers to degrees Celsius; "N" refers to normal or normality; "M" refers to molar or molarity; "mol" refers to mole or moles; "mmol" refers to millimole or millimoles; "kg" refers to kilogram or kilograms; "g" refers to gram or grams; "mg" refers to milligram or milligrams; "mL" refers
20 milliliter or milliliters; "L" refers to liter or liters; "bp" refers to boiling point; "mp" refers to melting point; "brine" refers to a saturated aqueous sodium chloride solution; etc.

25

EXAMPLE 1

Tomoxetine, (R)-N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine hydrochloride

Combine N-methyl-3-hydroxy-3-phenylpropylamine (10 g, 60.53 mmol) and potassium t-butoxide (7.5 g, 66.58 mmol) in
30 1,3-dimethyl-2-imidazolidinone (25 mL). Heat to dissolve, if necessary, before adding 2-fluorotoluene (20 mL, 181.6 mmol). Heat to about 110°C. After 20 hours, cool to ambient temperature and carefully add water and then toluene. Separate the layers and extract the organic layer
35 with water. Evaporate by distillation to reduce the volume of the organic layer to about 10 mL, cool to about 40° to 50°C, dilute with about 4.5 volumes of ethyl acetate (about

-15-

45 mL), and add (S)-(+)-mandelic acid (5.52 g, 36.3 mmol). Hold the temperature at about 40° to 45°C and seed with (R)-N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine (S)-(+)-mandelic acid salt. After about 2 hours, cool to about 0°
5 to 5°C to give a solid. Collect the solid by filtration, rinse with ethyl acetate and dry *in vacuo* at about 80°C. Combine (R)-N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine (S)-(+)-mandelic acid salt (4.36 g, 10.7 mmol), water (13.5 mL), aqueous 50% sodium hydroxide solution (0.99 g, 12.32
10 mmol), and methyl t-butyl ether (14 mL). Heat to about 40° to 45°C and stir until the solids dissolve. Separate the layers and extract the organic layer with water. Dilute the organic layer with methyl t-butyl ether (about 17 mL) and dry by distillation (collecting the azeotrope). Add an
15 additional amount of methyl t-butyl ether (about 17 mL) to bring the weight of the reaction mixture up to about 20.6 g. Add isopropanol (7.5 g) before slowly adding concentrated aqueous hydrochloric acid solution (1.11 g, 11.25 mmol) over about 3 hours. Stir the reaction mixture, cooling if
20 necessary to give a solid. Collect the solid by filtration, rinse with methyl t-butyl ether, and dry *in vacuo* at about 70° to 80°C to give the title compound.

EXAMPLE 2

25 Tomoxetine, (R)-N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine hydrochloride

Combine N-methyl-3-hydroxy-3-phenylpropylamine (10 g, 60.53 mmol) and potassium t-butoxide (7.5 g, 66.58 mmol) in 1,3-dimethyl-2-imidazolidinone (25 mL). Heat to dissolve,
30 if necessary, before adding 2-fluorotoluene (20 mL, 181.6 mmol). Heat to about 110°C. After 20 hours, cool to ambient temperature and carefully add water and then toluene. Separate the layers and extract the organic layer with water. Evaporate by distillation to reduce the volume
35 of the organic layer until about 3 g of toluene remains. Add toluene (about 12.5 g), cool to about 40° to 50°C, dilute with ethyl acetate (about 45 g), and add (S)-(+)-


-16-

mandelic acid (5.52 g, 36.3 mmol). Hold the temperature at about 40° to 45°C and seed with (R)-N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine (S)-(+)-mandelic acid salt. After about 2 hours, cool to about 0° to 5°C to give
5 a solid. Collect the solid by filtration, rinse with ethyl acetate and dry in vacuo at about 80°C. Combine (R)-N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine (S)-(+)-mandelic acid salt (4.36 g, 10.7 mmol), water (13.5 mL),
10 aqueous 50% sodium hydroxide solution (0.99 g, 12.32 mmol), and methyl t-butyl ether (14 mL). Heat to about 40° to 45°C and stir until the solids dissolve. Separate the layers and extract the organic layer with water. Dilute the organic layer with methyl t-butyl ether (about 17 mL) and dry by distillation (collecting the azeotrope). Add an additional
15 amount of methyl t-butyl ether (about 17 mL) to bring the weight of the reaction mixture up to about 20.6 g. Add isopropanol (7.5 g) before slowly adding concentrated aqueous hydrochloric acid solution (1.11 g, 11.25 mmol) over about 3 hours. Stir the reaction mixture, cooling if
20 necessary to give a solid. Collect the solid by filtration, rinse with methyl t-butyl ether, and dry in vacuo at about 70° to 80°C to give the title compound.

-17-

WE CLAIM:

- 5 1. A process for preparing 3-aryloxy-3-arylpropylamines of the formula



10 wherein


Ar is phenyl or 2-thienyl,

Ar₁ is 1-naphthyl, 2-methoxyphenyl, 2-thiomethoxyphenyl, or 2-methylphenyl;

G is hydrogen or methyl,

- 15 and the pharmaceutically-acceptable addition salts thereof comprising the steps of:

(a) reacting an alkoxide of a 3-hydroxy-3-arylpropylamine of the formula



20

wherein

Ar and G are as defined above

with a haloaromatic of the formula

25 Ar_1-X

wherein

Ar₁-X is 2-fluorotoluene, 2-chlorotoluene, 1-

fluoronaphthalene, 1-chloronaphthalene, 2-fluoroanisole, 2-chloroanisole, 2-fluorothioanisole, or 2-chlorothioanisole,

- 30 in 1,3-dimethyl-2-imidazolidinone or N-methylpyrrolidinone to give the 3-aryloxy-3-arylpropylamine;

-18-

(b) N-demethylated of the 3-aryloxy-3-arylpropylamine wherein G is methyl to give the 3-aryloxy-3-arylpropylamine wherein G is hydrogen;

(c) optional resolution of the 3-aryloxy-3-arylpropylamine to give a specific isomer of the 3-aryloxy-3-arylpropylamine; and

(d) optional formation of an acid addition salt using a pharmaceutically-acceptable acid.

2. A process according to Claim 1 wherein 1,3-dimethyl-2-imidazolidinone is used.

3. A process according to Claim 2 wherein the 3-hydroxy-3-arylpropylamine is (S)-N,N-dimethyl-3-(2-thienyl)-3-hydroxypropylamine and the haloaromatic is 1-fluoronaphthalene.

4. A process for preparing tomoxetine and the pharmaceutically-acceptable addition salts thereof comprising the steps of:

(a) reacting an alkoxide of N-methyl-3-phenyl-3-hydroxypropylamine with 2-fluorotoluene in 1,3-dimethyl-2-imidazolidinone to give N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine;

(b) resolution of N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine to give (R)-N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine (tomoxetine); and

(c) optional formation of an acid addition salt using a pharmaceutically-acceptable acid.

5. A process for preparing tomoxetine and the pharmaceutically-acceptable addition salts thereof comprising the steps of:

(a) reacting an alkoxide of N,N-dimethyl-3-phenyl-3-hydroxypropylamine) with 2-fluorotoluene in 1,3-dimethyl-2-imidazolidinone to give N,N-dimethyl-3-(2-methylphenoxy)-3-phenylpropylamine;

-19-

(b) N-demethylated N,N-dimethyl-3-(2-methylphenoxy)-3-phenylpropylamine to give N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine;

(c) resolution of N-methyl-3-(2-methylphenoxy)-3-

5 phenylpropylamine to give (R)-N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine; and

(d) optional formation of an acid addition salt using a pharmaceutically-acceptable acid.

10 6. A process for preparing N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine comprising the steps of:

(a) reacting an alkoxide of N-methyl-3-phenyl-3-hydroxypropylamine with 2-fluorotoluene in 1,3-dimethyl-2-imidazolidinone.

15


7. A process for preparing N-methyl-3-(2-methylphenoxy)-3-phenylpropylamine comprising the steps of:

(a) reacting an alkoxide of N,N-dimethyl-3-phenyl-3-hydroxypropylamine) with 2-fluorotoluene in 1,3-dimethyl-2-imidazolidinone to give N,N-dimethyl-3-(2-methylphenoxy)-3-phenylpropylamine;

20

(b) N-demethylated N,N-dimethyl-3-(2-methylphenoxy)-3-phenylpropylamine.

25 8. A process for preparing 3-aryloxy-3-arylpropylamines of the formula



30 wherein


Ar is phenyl or 2-thienyl,

Ar₁ is 1-naphthyl, 2-methoxyphenyl, 2-thiomethoxyphenyl, or 2-methylphenyl;

-20-

G is hydrogen or methyl,
and the pharmaceutically acceptable salts thereof, which
comprises reacting an alkoxide of N-methyl-3-hydroxy-3-
arylpropylamine of the formula

5



or an N-protected derivative thereof,

Ar and G are as defined above with an unactivated
haloaromatic, characterized in that, 1,3-dimethyl-2-
imidazolidinone or N-methylpyrrolidinone is used as solvent.

9. A process for preparing tomoxetine and the
pharmaceutically-acceptable addition salts thereof, which
comprises, reacting an alkoxide of N-methyl-3-phenyl-3-
hydroxypropylamine, or an N-protected derivative thereof,
with 2-fluorotoluene, characterized in that, 1,3-dimethyl-2-
imidazolidinone is used as solvent.

20

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/06423

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07C217/48 C07C213/06 C07B57/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07C C07B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 362 886 A (BERGLUND RICHARD A) 8 November 1994 (1994-11-08) cited in the application column 1, line 60 -column 3, line 2; especially column 1 line 67 -column 2 line 2 example 1	1-9
X	US 5 166 437 A (KAIRISALO PEKKA J ET AL) 24 November 1992 (1992-11-24) cited in the application example 2	1-9
X	US 5 847 214 A (ROSSETTI VITTORIO ET AL) 8 December 1998 (1998-12-08) column 3, line 38 - line 53 examples 1,3	1-9
	-/-	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

12 July 2000

Date of mailing of the international search report

19/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel.: (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

O'Sullivan, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/06423

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>ES 2 120 368 A (ALMIRALL PRODESFARMA SA) 16 October 1998 (1998-10-16) column 3, line 35 - line 48</p>	1-9

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/06423

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5362886	A	08-11-1994	AU 685494 B	22-01-1998
			AU 7572094 A	04-05-1995
			BR 9404045 A	13-06-1995
			CA 2133899 A	13-04-1995
			CN 1109470 A	04-10-1995
			CZ 9402465 A	17-05-1995
			EP 0650965 A	03-05-1995
			FI 944773 A	13-04-1995
			HU 68943 A	28-08-1995
			IL 111188 A	15-06-1998
			JP 7188065 A	25-07-1995
			NO 943825 A	18-04-1995
			NZ 264633 A	22-09-1997
			PL 305326 A	18-04-1995
			RU 2127269 C	10-03-1999
			US 5491243 A	13-02-1996
			ZA 9407839 A	09-04-1996
US 5166437	A	24-11-1992	FI 81083 B	31-05-1990
			AT 100440 T	15-02-1994
			AU 620497 B	20-02-1992
			AU 4996990 A	06-09-1990
			BG 51156 A	15-02-1993
			CA 2010410 A,C	03-09-1990
			CS 9001003 A	15-09-1991
			DD 297806 A	23-01-1992
			DE 69006072 D	03-03-1994
			DE 69006072 T	30-06-1994
			DK 49790 A	04-09-1990
			DK 391070 T	28-02-1994
			EP 0391070 A	10-10-1990
			ES 2062131 T	16-12-1994
			HR 921278 A	31-08-1995
			HU 54619 A,B	28-03-1991
			IE 63650 B	31-05-1995
			IL 92802 A	25-01-1994
			JP 2062299 C	24-06-1996
			JP 2282352 A	19-11-1990
			JP 7091244 B	04-10-1995
			NO 901013 A,B,	04-09-1990
			PL 163884 B	31-05-1994
			PT 93302 A	07-11-1990
			RO 105802 A	30-12-1992
			SI 9010015 A,B	30-06-1998
			SU 1779244 A	30-11-1992
			YU 1590 A	31-10-1991
			ZA 9000739 A	28-11-1990
US 5847214	A	08-12-1998	IT MI961438 A	12-01-1998
ES 2120368	A	16-10-1998	NONE	